

HIV Prevention and Transmission Myths among a Sample of Heterosexually Active Adults in South Florida

**2009 National HIV Prevention Conference
Atlanta, GA, August 24, 2009**

Dano Beck, M.S.W., Lisa Metsch, Ph.D.,
Marlene LaLota, M.P.H., Gabriel A. Cardenas, M.P.H.,
David Forrest, Ph.D., Spencer Lieb, M.P.H.




Background

- Estimates of annual HIV incidence in the U.S. emphasize the need for continued and effective prevention activities
- It is important that future prevention efforts consider the full range of prevention and transmission beliefs among persons at risk for or living with HIV
- Holding false or conspiracy beliefs about HIV and its transmission has been associated with behaviors that can lead to increased HIV transmission

National HIV Behavioral Surveillance (NHBS)

- **CDC-funded study in over 20 U.S. MSAs with high AIDS prevalence**
 - FL NHBS sites: Miami and Fort Lauderdale
- **Collects cross-sectional data**
 - HIV prevalence, risk behaviors, testing behaviors, and use of prevention services
- **Study cycles rotate by population**
 - MSM, IDUs, and at-risk heterosexuals



Behavioral Surveillance Among Heterosexuals At Risk for HIV Infection (NHBS-HET)


- Conducted in 2007
- Heterosexual at risk for HIV infection: an adult male or female with
 - a physical or social connection to a “high- risk area” and
 - at least one opposite-sex partner in the past year
- High-risk areas: census tracts with high rates of poverty and heterosexually-acquired HIV/AIDS cases
- Venue-based sampling was used to recruit 1,224 eligible participants within selected high-risk areas
- Measures on HIV prevention and transmission beliefs were included in Florida’s local questionnaire

NHBS-HET Eligibility Criteria

- 18-50 years of age
- Lives in Miami-Dade, Broward, or Palm Beach county
- Male or female (not transgender)
- Reports vaginal or anal sex with a person of the opposite sex in the past 12 months
- Able to complete interview in English or Spanish
- Not a previous NHBS-HET participant

Demographic Characteristics of Sample N=1,224

- Mean age: 35 years
- 58% male, 42% female
- Majority black



*Results are preliminary


Characteristics of Sample (cont'd)

- 91% self-identify as heterosexual, 9% as bisexual
- Majority (85%) born in U.S.
- 2/3 (63%) without health insurance at time of interview
- Majority (60%) have an annual income under \$10,000
- Half (50%) unemployed at time of interview
- 42% depressed (according to CES-D 10)


Characteristics of Sample (cont'd)

- 1/3 (33%) incarcerated in past 12 months
- 1/4 (28%) used cocaine or crack in past 12 months
- 1/5 (21%) self-reported being hepatitis C+
- 13% homeless in past 12 months
- 1 in 12 (8%) tested HIV+, and 1 in 24 (4%) had a previously undiagnosed infection


Beliefs on How HIV Can Be Spread, N=1,221



Beliefs on How HIV Can Be Prevented, N=1,221



Beliefs on How HIV Can Be Prevented (cont'd), N=1,221



Odds Ratio Estimates for Having One or More Incorrect Prevention Beliefs

Logistic Regression

	<u>Adj OR</u>	<u>95% LCL</u>	<u>95%UCL</u>
Female ^x	0.77	0.58	1.02
Age ^x	1.01	1.0	1.03
Hispanic ^{*x}	3.35	1.19	9.48
Black (non-Hispanic) ^{*x}	2.02	0.74	5.50
Positive HIV Status ^{**}	0.58	0.27	1.25
Unknown HIV Status ^{**}	1.55	1.16	2.07
Depressed	2.14	1.63	2.82

x Forced into model

*Reference for race/ethnicity is white (non-Hispanic)

**Reference for self-reported HIV status is negative

Odds Ratio Estimates for Having One or More Incorrect Transmission Beliefs

Logistic Regression

	<u>Adj OR</u>	<u>95% LCL</u>	<u>95%UCL</u>
Female ^x	0.87	0.66	1.13
Miami ^x	0.83	0.64	1.07
Age ^x	0.98	0.97	0.99
Hispanic ^{*x}	1.20	0.47	3.05
Black (non-Hispanic) ^{*x}	1.46	0.60	3.56
Positive HIV Status ^{**}	0.50	0.23	1.08
Unknown HIV Status ^{**}	1.49	1.12	1.97
Depressed	1.32	1.01	1.72
Heavy Alcohol Use	1.78	1.33	2.38
Visited HIV Provider in Past 12 Months	0.73	0.56	0.96

x Forced into model.

*Reference for race/ethnicity is white (non-Hispanic)

**Reference for self-reported HIV status is negative

Limitations

- Data are preliminary
- Data are predominantly self-reported and subject to recall bias
- Data are cross sectional and time order cannot be established
- Findings can only be generalized to the population meeting eligibility criteria who attended venues on the sampling frame

Implications

- While most (>90%) participants held correct beliefs about HIV prevention and transmission, over 1/3 reported one or more misconceptions
- Participants reporting myths were commonly unaware of their HIV status and may be at increased risk for transmitting HIV to others
- Future prevention efforts should debunk myths about HIV prevention and transmission
- Service providers should link clients to HIV testing, mental health services, and other healthcare services

THANK YOU!

**A BIG THANK YOU TO THE FLORIDA
NHBS-HET TEAM FOR THEIR
OUTSTANDING WORK!**

David Forrest, PhD

Emelina Martinez

Roxana Bell

Cristobal Plaza

Gabriel A. Cardenas, MPH

Wei Zhao, MD

Cheryl Riles

Lottie Watson

Dwight Lavant

Contact Information

Dano Beck, M.S.W.
Behavioral Surveillance
Coordinator
Bureau of HIV/AIDS
Florida Department of Health

Dano_Beck@doh.state.fl.us

David Forrest, Ph.D.
Field Operations Director
Dept. of Epidemiology &
Public Health
University of Miami

DForrest@med.miami.edu

